


GalNAcEXO™

FOR RESEARCH
USE ONLY
www.genovis.com

STORE AT
-20°C



SmartEnzymes™



GENOVIS

INSTRUCTIONS FOR PRODUCT

GalNAcEXO 2000 units (G1-NA1-020)

Digestion of up to 2 mg glycoprotein

1 Prepare GalNAcEXO

- Reconstitute GalNAcEXO in 100 μ l ddH₂O to a concentration of 20 units/ μ l



2 Add GalNAcEXO

- Add 1 unit GalNAcEXO / 1 μ g glycoprotein



3 Digestion

- Incubate for 2 h to 18 h at 37°C



PRODUCT DESCRIPTION

GalNAcEXO is an exo- α -N-Acetylgalactosaminidase for efficient hydrolysis of α -N-acetylgalactosamine (GalNAc) linked to serine or threonine residues in glycoproteins (Tn antigen). GalNAcEXO has some activity on other terminal α -linked GalNAc. GalNAcEXO hydrolyzes glycoproteins under native conditions and is highly active in a pH range from 6.0 to 7.6. No cofactors or special buffers are required. The enzyme in GalNAcEXO is derived from *Akkermansia muciniphila*, expressed in *E. coli* with a His-tag, and has a molecular weight of 52kDa.

Unit Definition

One unit of GalNAcEXO catalyzes the hydrolysis of α -linked GalNAc residues from >95% of 2 nmol 4-Nitrophenyl 2-acetamido-2-deoxy- α -D-galactopyranoside when incubated in 20mM Tris, 1% EtOH pH 6.8 at 37°C for 30 min.

Content and Storage

GalNAcEXO is supplied lyophilized in TBS pH 7.6, with no preservatives added.

GalNAcEXO is shipped cold and should be stored at -20°C upon arrival.

After reconstitution GalNAcEXO is stable for 1 month at $+4-8^{\circ}\text{C}$.

GalNAcEXO is for R&D use only.

DETAILED PROTOCOL

Additional Materials Required

- Reaction buffer¹: 20 mM Tris pH 6.8

Sample Preparation

- Prepare the glycoprotein of interest in the reaction buffer at a concentration of 0.5-5.0 mg/ml.

Digestion of Glycoproteins

1 Prepare GalNAcEXO

- Reconstitute GalNAcEXO in 100 μ l ddH₂O to a concentration of 20 units/ μ l

2 Add GalNAcEXO

- Add 1 unit GalNAcEXO / 1 μ g glycoprotein²

3 Digestion

- Incubate for 2 to 18 h at 37°C

Optimization of enzyme concentrations and incubation time may be needed depending on the substrate.

Notes

1. GalNAcEXO displays high activity in buffers with pH values from 6.0 to 7.6 and over a wide range of ionic strength. Some optimizations might be required if a buffer other than the recommended reaction buffer is used.
2. A higher enzyme concentration may increase digestion efficiency of individual glycoproteins. This requires optimization.

Quality Control

GalNAcEXO is tested to meet the specifications and lot-to-lot consistency.

GalNAcEXO is tested for the absence of microbial contamination using blood agar plates, Sabouraud dextrose agar plates and fluid thioglycollate medium.

Related Products

SialEXO®

For complete removal of α 2-3, α 2-6 and α 2-8 linked sialic acids.

GalactEXO™

GalNAcEXO for complete removal of both β 1-3 and β 1-4 linked galactoses.

OglyZOR®

Specific hydrolysis of core 1 O-glycan disaccharides on native glycoproteins.

Legal and Disclaimers

All rights reserved. Genovis products may be covered by one or more patents, trademarks and copyrights owned or controlled by Genovis AB.

For more information about commercial rights, please contact Genovis team at info@genovis.com

Genovis products are intended for research use only. They are not intended to be used for therapeutic or diagnostic purposes in humans or animals.

© Genovis 2020



USA & Canada

Genovis Inc.
245 First Street, Suite 1800
Cambridge, MA 02142
USA

Customer service: 1-617-444-8421
Order phone (toll free): 1-855-782-0084
Order fax: 1-858-524-3006
Email: orders.us@genovis.com

EMEA & Asia

Genovis AB
Box 790
SE-220 07 Lund
Sweden

Customer service: 0046 (0)46 10 12 30
Order phone: 0046 (0)46 10 12 30
Order fax: 0046 (0)46 12 80 20
Email: order@genovis.com